

Notes on Graph Product Structure Theory

Zdeněk Dvořák, Tony Huynh, Gwenaël Joret, Chun-Hung Liu, and David R. Wood

Abstract It was recently proved that every planar graph is a subgraph of the strong product of a path and a graph with bounded treewidth. This paper surveys generalisations of this result for graphs on surfaces, minor-closed classes, various non-minor-closed classes, and graph classes with polynomial growth. We then explore how graph product structure might be applicable to more broadly defined graph classes. In particular, we characterise when a graph class defined by a cartesian or strong product has bounded or polynomial expansion. We then explore graph product structure theorems for various geometrically defined graph classes, and present several open problems.

Zdeněk Dvořák
Charles University, Prague, Czech Republic
Supported by project 17-04611S (Ramsey-like aspects of graph coloring) of Czech Science Foundation
e-mail: rakdver@iuuk.mff.cuni.cz

Tony Huynh and David R. Wood
School of Mathematics, Monash University, Melbourne, Australia
Research supported by the Australian Research Council
e-mail: tony.bourbaki@gmail.com, david.wood@monash.edu

Gwenaël Joret
Département d'Informatique, Université libre de Bruxelles, Brussels, Belgium
Research supported by the Wallonia-Brussels Federation of Belgium, and by the Australian Research Council
e-mail: gjoret@ulb.ac.be

Chun-Hung Liu
Department of Mathematics, Texas A&M University, College Station, Texas, USA
Partially supported by the NSF under Grant No. DMS-1929851
e-mail: chliu@math.tamu.edu

1 Introduction

Studying the structure of graphs is a fundamental topic of broad interest in combinatorial mathematics. At the forefront of this study is the Graph Minor Theorem of Robertson and Seymour [47], described by Diestel [8] as “among the deepest theorems mathematics has to offer”. At the heart of the proof of this theorem is the Graph Minor Structure Theorem, which shows that any graph in a minor-closed family¹ can be constructed using four ingredients: graphs on surfaces, vortices, apex vertices, and clique-sums. Graphs of bounded genus, and in particular planar graphs are basic building blocks in graph minor structure theory. Indeed, the theory says nothing about the structure of planar graphs. So it is natural to ask whether planar graphs can be described in terms of some simpler graph classes. In a recent breakthrough, Dujmović, Joret, Micek, Morin, Ueckerdt, and Wood [15, 16] provided an answer to this question by showing that every planar graph is a subgraph of the strong product² of a graph of bounded treewidth³ and a path.

Theorem 1 ([15, 16]). *Every planar graph is a subgraph of:*

- (a) $H \boxtimes P$ for some graph H of treewidth at most 8 and for some path P ;
- (b) $H \boxtimes P \boxtimes K_3$ for some graph H of treewidth at most 3 and for some path P .

This *graph product structure theorem* is attractive since it describes planar graphs in terms of graphs of bounded treewidth, which are considered much simpler than planar graphs. For example, many NP-complete problem remain NP-complete on planar graphs but are polynomial-time solvable on graphs of bounded treewidth.

Despite being only 10 months old, Theorem 1 is already having significant impact. Indeed, it has been used to solve two major open problems and make additional progress on two other longstanding problems:

- Dujmović et al. [15, 16] use Theorem 1 to show that planar graphs have queue layouts with a bounded number of queues, solving a 27 year old problem of Heath, Leighton, and Rosenberg [33].

¹ A graph H is a *minor* of a graph G if a graph isomorphic to H can be obtained from a subgraph of G by contracting edges. A class of graphs \mathcal{G} is *minor-closed* if for every graph $G \in \mathcal{G}$ every minor of G is in \mathcal{G} , and some graph is not in \mathcal{G} . A graph G is *H -minor-free* if H is not a minor of G .

² The *cartesian product* of graphs A and B , denoted by $A \square B$, is the graph with vertex set $V(A) \times V(B)$, where distinct vertices $(v, x), (w, y) \in V(A) \times V(B)$ are adjacent if: $v = w$ and $xy \in E(B)$; or $x = y$ and $vw \in E(A)$. The *strong product* of graphs A and B , denoted by $A \boxtimes B$, is the graph with vertex set $V(A) \times V(B)$, where distinct vertices $(v, x), (w, y) \in V(A) \times V(B)$ are adjacent if: $v = w$ and $xy \in E(B)$; or $x = y$ and $vw \in E(A)$; or $vw \in E(A)$ and $xy \in E(B)$. If X is a subgraph of $A \square B$, then the *projection* of X into A is the set of vertices $v \in V(A)$ such that $(v, w) \in V(X)$ for some $w \in V(B)$.

³ A *tree decomposition* of a graph G is a collection $(B_x \subseteq V(G) : x \in V(T))$ of subsets of $V(G)$ (called *bags*) indexed by the nodes of a tree T , such that (i) for every edge $uv \in E(G)$, some bag B_x contains both u and v , and (ii) for every vertex $v \in V(G)$, the set $\{x \in V(T) : v \in B_x\}$ induces a non-empty (connected) subtree of T . The *width* of a tree decomposition is the size of the largest bag minus 1. The *treewidth* of a graph G , denoted by $\text{tw}(G)$, is the minimum width of a tree decomposition of G . See [3, 4, 32, 45, 46] for surveys on treewidth. A *path decomposition* is a tree decomposition where the underlying tree is a path. The *pathwidth* of a graph G , denoted by $\text{pw}(G)$, is the minimum width of a path decomposition of G .

- Dujmović, Eppstein, Joret, Morin, and Wood [12] use Theorem 1 to show that planar graphs can be nonrepetitively coloured with a bounded number of colours, solving a 17 year old problem of Alon, Grytczuk, Hałuszczak, and Riordan [1].
- Dębski, Felsner, Micek, and Schröder [10] use Theorem 1 to prove the best known results on p -centred colourings of planar graphs, reducing the bound from $O(p^{19})$ to $O(p^3 \log p)$.
- Bonamy, Gavoille, and Pilipczuk [5] use Theorem 1 to give more compact graph encodings of planar graphs. In graph-theoretic terms, this implies the existence of a graph with $n^{4/3+o(1)}$ vertices that contains each planar graph with at most n vertices as an induced subgraph. This work improves a sequence of results that goes back 27 years to the introduction of implicit labelling schemes by Kannan, Naor, and Rudich [34].

The first goal of this paper is to introduce several product structure theorems that have been recently established, most of which generalise Theorem 1. First Section 2 considers minor-closed classes. Then Section 3 considers several examples of non-minor-closed classes. Section 4 introduces the notion of graph classes with polynomial growth and their characterisation in terms of strong products of paths due to Krauthgamer and Lee [36]. We prove an extension of this result for strong products of graphs of given pathwidth.

The remaining sections explore how graph product structure might be applicable to more broadly defined graph classes. The following definition by Nešetřil and Ossona de Mendez [40] provides a setting for this study⁴. A graph class \mathcal{G} has *bounded expansion* with *expansion function* $f: \mathbb{Z}^+ \rightarrow \mathbb{R}$ if, for every graph $G \in \mathcal{G}$ and for all disjoint subgraphs B_1, \dots, B_t of radius at most r in G , every subgraph of the graph obtained from G by contracting each B_i into a vertex has average degree at most $f(r)$. When $f(r)$ is a constant, \mathcal{G} is contained in a proper minor-closed class. As $f(r)$ is allowed to grow with r we obtain larger and larger graph classes. A graph class \mathcal{G} has *linear expansion* if \mathcal{G} has bounded expansion with an expansion function in $O(r)$. A graph class \mathcal{G} has *polynomial expansion* if \mathcal{G} has bounded expansion with an expansion function in $O(r^c)$, for some constant c .

We characterise when a graph class defined by a cartesian or strong product has bounded or polynomial expansion. For $\star \in \{\boxtimes, \square\}$ and for hereditary⁵ graph classes \mathcal{G}_1 and \mathcal{G}_2 , let

$$\mathcal{G}_1 \star \mathcal{G}_2 := \{G : G \subseteq G_1 \star G_2, G_1 \in \mathcal{G}_1, G_2 \in \mathcal{G}_2\}.$$

Note that $\mathcal{G}_1 \star \mathcal{G}_2$ is hereditary. Sections 5 and 6 characterise when $\mathcal{G}_1 \star \mathcal{G}_2$ has bounded or polynomial expansion. In related work, Wood [53] characterised when

⁴ Let $d_G(u, v)$ be the distance between vertices u and v in a graph G . For a vertex v in a graph G and $r \in \mathbb{N}$, let $N_G^r(v)$ be the set of vertices of G at distance exactly r from v , and let $N_G[v]$ be the set of vertices at distance at most r from v . The set $N_G^r[v]$ is called an *r-ball*. We drop the subscript G when the graph is clear from the context.

⁵ A class of graphs is *hereditary* if it is closed under induced subgraphs.

$\mathcal{G}_1 \square \mathcal{G}_2$ has bounded Hadwiger number, and Pecaninovic [43] characterised when $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has bounded Hadwiger number.

Section 7 explores graph product structure theorems for various geometrically defined graph classes. We show that multi-dimensional unit-disc graphs have a product structure theorem, and discusses whether two other naturally defined graph classes might have product structure theorems. We finish with a number of open problems in Section 8.

2 Minor-Closed Classes

Here we survey results generalising Theorem 1 for minor-closed classes. First consider graphs embeddable on a fixed surface⁶.

Theorem 2 ([15, 16]). *Every graph of Euler genus g is a subgraph of:*

- (a) $H \boxtimes P \boxtimes K_{\max\{2g,1\}}$ for some graph H of treewidth at most 9 and for some path P ;
- (b) $H \boxtimes P \boxtimes K_{\max\{2g,3\}}$ for some graph H of treewidth at most 4 and for some path P .
- (c) $(K_{2g} + H) \boxtimes P$ for some graph H of treewidth at most 8 and some path P .

Here $A + B$ is the complete join of graphs A and B . The proof of Theorem 2 uses an elegant cutting lemma to reduce to the planar case.

Theorem 2 is generalised as follows. A graph X is *apex* if $X - v$ is planar for some vertex v .

Theorem 3 ([15, 16]). *For every apex graph X , there exists $c \in \mathbb{N}$ such that every X -minor-free graph G is a subgraph of $H \boxtimes P$ for some graph H of treewidth at most c and some path P .*

The proof of Theorem 3 is based on the Graph Minor Structure Theorem of Robertson and Seymour [49] and in particular a strengthening of it by Dvořák and Thomas [21].

For an arbitrary proper minor-closed class, apex vertices are unavoidable; in this case Dujmović et al. [15, 16] proved the following product structure theorem.

Theorem 4 ([15, 16]). *For every proper minor-closed class \mathcal{G} there exist $k, a \in \mathbb{N}$ such that every graph $G \in \mathcal{G}$ can be obtained by clique-sums of graphs G_1, \dots, G_n such that for $i \in \{1, \dots, n\}$,*

$$G_i \subseteq (H_i \boxtimes P_i) + K_a,$$

for some graph H_i with treewidth at most k and some path P_i .

⁶ The *Euler genus* of the orientable surface with h handles is $2h$. The *Euler genus* of the non-orientable surface with c cross-caps is c . The *Euler genus* of a graph G is the minimum Euler genus of a surface in which G embeds (with no crossings). See [39] for background on embeddings of graphs on surfaces.

If we assume bounded maximum degree, then apex vertices in the Graph Minor Structure Theorem can be avoided, which leads to the following theorem of Dujmović, Esperet, Morin, Walczak, and Wood [14].

Theorem 5 ([14]). *For every proper minor-closed class \mathcal{G} , every graph in \mathcal{G} with maximum degree Δ is a subgraph of $H \boxtimes P$ for some graph H of treewidth $O(\Delta)$ and for some path P .*

It is worth highlighting the similarity of Theorem 5 and the following result of Ding and Oporowski [9] (refined in [52]). Theorem 6 says that graphs of bounded treewidth and bounded degree are subgraphs of the product of a tree and a complete graph of bounded size, whereas Theorem 5 says that graphs excluding a minor and with bounded degree are subgraphs of the product of a bounded treewidth graph and a path.

Theorem 6 ([9, 52]). *Every graph with maximum degree $\Delta \geq 1$ and treewidth at most $k \geq 1$ is a subgraph of $T \boxtimes K_{18k\Delta}$ for some tree T .*

3 Non-Minor Closed Classes

A recent direction pursued by Dujmović, Morin, and Wood [17] studies graph product structure theorems for various non-minor-closed graph classes. First consider graphs that can be drawn on a surface of bounded genus and with a bounded number of crossings per edge. A graph is (g, k) -planar if it has a drawing in a surface of Euler genus at most g such that each edge is involved in at most k crossings. Even in the simplest case, there are $(0, 1)$ -planar graphs that contain arbitrarily large complete graph minors [13].

Theorem 7 ([17]). *Every (g, k) -planar graph is a subgraph of $H \boxtimes P$, for some graph H of treewidth $O(gk^6)$ and for some path P .*

Map and string graphs provide further examples of non-minor-closed classes that have product structure theorems.

Map graphs are defined as follows. Start with a graph G_0 embedded in a surface of Euler genus g , with each face labelled a ‘nation’ or a ‘lake’, where each vertex of G_0 is incident with at most d nations. Let G be the graph whose vertices are the nations of G_0 , where two vertices are adjacent in G if the corresponding faces in G_0 share a vertex. Then G is called a (g, d) -map graph. A $(0, d)$ -map graph is called a (plane) d -map graph; see [7, 26] for example. The $(g, 3)$ -map graphs are precisely the graphs of Euler genus at most g ; see [13]. So (g, d) -map graphs generalise graphs embedded in a surface, and we now assume that $d \geq 4$ for the remainder of this section.

Theorem 8 ([17]). *Every (g, d) -map graph is a subgraph of:*

- $H \boxtimes P \boxtimes K_{O(d^2g)}$, where H is a graph with treewidth at most 14 and P is a path,
- $H \boxtimes P$, where H is a graph with treewidth $O(gd^2)$ and P is a path.

A *string graph* is the intersection graph of a set of curves in the plane with no three curves meeting at a single point; see [27, 28, 42] for example. For $\delta \in \mathbb{N}$, if each curve is in at most δ intersections with other curves, then the corresponding string graph is called a δ -*string graph*. A (g, δ) -*string graph* is defined analogously for curves on a surface of Euler genus at most g .

Theorem 9 ([17]). *Every (g, δ) -string graph is a subgraph of $H \boxtimes P$, for some graph H of treewidth $O(g\delta^7)$ and some path P .*

Theorems 7 to 9 all follow from a more general result of Dujmović et al. [17]. A collection \mathcal{P} of paths in a graph G is a (k, d) -*shortcut system* (for G) if:

- every path in \mathcal{P} has length at most k , and
- for every $v \in V(G)$, the number of paths in \mathcal{P} that use v as an internal vertex is at most d .

Each path $P \in \mathcal{P}$ is called a *shortcut*; if P has endpoints v and w then it is a vw -*shortcut*. Given a graph G and a (k, d) -shortcut system \mathcal{P} for G , let $G^{\mathcal{P}}$ denote the supergraph of G obtained by adding the edge vw for each vw -shortcut in \mathcal{P} .

Theorem 10 ([17]). *Let G be a subgraph of $H \boxtimes P$, for some graph H of treewidth at most t and for some path P . Let \mathcal{P} be a (k, d) -shortcut system for G . Then $G^{\mathcal{P}}$ is a subgraph of $J \boxtimes P'$ for some graph J of treewidth at most $d(k^3 + 3k) \binom{k+t}{t} - 1$ and some path P' .*

Theorems 7 to 9 are then proved by simply constructing a shortcut system. For example, by adding a dummy vertex at each crossing, Dujmović et al. [17] noted that every (g, k) -planar graph is a subgraph of $G^{\mathcal{P}}$ for some graph G of Euler genus at most g and for some $(k+1, 2)$ -shortcut system \mathcal{P} for G .

Powers of graphs can also be described by a shortcut system. The k -*th power* of a graph G is the graph G^k with vertex set $V(G^k) := V(G)$, where $vw \in E(G^k)$ if and only if $d_G(v, w) \leq k$. Dujmović et al. [17] noted that if a graph G has maximum degree Δ , then $G^k = G^{\mathcal{P}}$ for some $(k, 2k\Delta^k)$ -shortcut system \mathcal{P} . Theorem 10 then implies:

Theorem 11 ([17]). *For every graph G of Euler genus g and maximum degree Δ , the k -th power G^k is a subgraph of $H \boxtimes P$, for some graph H of treewidth $O(g\Delta^k k^8)$ and some path P .*

4 Polynomial Growth

This section discusses graph classes with polynomial growth. A graph class \mathcal{G} has *polynomial growth* if for some constant c , for every graph $G \in \mathcal{G}$, for each $r \geq 2$ every r -ball in G has at most r^c vertices. For example, every r -ball in an $n \times n$ grid graph is contained in a $(2r+1) \times (2r+1)$ subgrid, which has size $(2r+1)^2$; therefore the class of grid graphs has polynomial growth. More generally, let \mathbb{Z}^d be the strong product of d infinite two-way paths. That is, $V(\mathbb{Z}^d) = \{(x_1, \dots, x_d) : x_1, \dots, x_d \in \mathbb{Z}\}$ where distinct vertices (x_1, \dots, x_d) and (y_1, \dots, y_d) are adjacent in

\mathbb{Z}^d if and only if $|x_i - y_i| \leq 1$ for each $i \in \{1, \dots, d\}$. Then every r -ball in \mathbb{Z}^d has size at most $(2r+1)^d$. Krauthgamer and Lee [36] characterised the graph classes with polynomial growth as the subgraphs of \mathbb{Z}^d .

Theorem 12 ([36]). *Let G be a graph such that for some constant c and for every integer $r \geq 2$, every r -ball in G has at most r^c vertices. Then $G \subseteq \mathbb{Z}^{O(c \log c)}$.*

We show that a seemingly weaker condition also characterises graph classes with polynomial growth. (We emphasise that in Theorem 13, H_1 does not necessarily have bounded maximum degree.)

Theorem 13. *The following are equivalent for a class of graphs \mathcal{G} :*

- (1) \mathcal{G} has polynomial growth,
- (2) there exists $d \in \mathbb{N}$ such that every graph in \mathcal{G} is a subgraph of \mathbb{Z}^d ,
- (3) there exist $d, k, \ell, \Delta \in \mathbb{N}$ such that for every graph $G \in \mathcal{G}$ there exist graphs H_1, \dots, H_d such that:
 - G has maximum degree Δ ,
 - $\text{pw}(H_i) \leq k$ for each $i \in \{1, \dots, k\}$,
 - H_i has maximum degree at most ℓ for each $i \in \{2, \dots, d\}$,
 - $G \subseteq H_1 \boxtimes H_2 \boxtimes \dots \boxtimes H_d$.

Proof. Krauthgamer and Lee [36] proved that (1) and (2) are equivalent. It is immediate that (2) implies (3) with $k = 1$ and $\ell = 2$ and $\Delta = 3^d - 1$. So it suffices to show that (3) implies (1). Consider graphs $G \in \mathcal{G}$ and H_1, \dots, H_d satisfying (3). For $i \in \{2, \dots, d\}$, by Lemma 14 below (with $d = 0$), every r -ball in H_i has at most $(1 + \ell)^k (2r + 1)^{k+1}$ vertices. By the result of Krauthgamer and Lee [36], $H_i \subseteq \mathbb{Z}^c$ for some $c = c(k, \ell)$. Thus

$$G \subseteq H_1 \boxtimes \mathbb{Z}^{c(d-1)}.$$

By Lemma 14 again, every r -ball in G has size at most

$$(1 + \Delta)^k (2r + 1)^{(k+1)(c(d-1)+1)},$$

which is at most $r^{c'}$ for some $c' = c'(c, \Delta, k)$ and $r \geq 2$. Hence (1) holds. \square

Lemma 14. *For every graph H with pathwidth at most $k \in \mathbb{N}_0$, for every connected subgraph G of $H \boxtimes \mathbb{Z}^d$ with radius at most r and maximum degree at most Δ ,*

$$|V(G)| \leq (1 + \Delta)^k (2r + 1)^{(k+1)(d+1)}.$$

Proof. The BFS spanning tree of G rooted at the centre of G has radius at most r . So it suffices to prove the result when G is a tree. We proceed by induction on $k \geq 0$ with the following hypothesis: For every graph H with pathwidth at most $k \in \mathbb{N}_0$, for every subtree T of $H \boxtimes \mathbb{Z}^d$ with radius at most r and maximum degree at most Δ ,

$$|V(T)| \leq (1 + \Delta)^k (2r + 1)^{(k+1)(d+1)}.$$

Since T is connected, we may assume that H is connected. Since T has radius at most r ,

$$T \subseteq H \boxtimes P_1 \boxtimes \cdots \boxtimes P_d,$$

where each P_i is a path on $2r + 1$ vertices.

In the base case $k = 0$, we have $H = K_1$ and $T \subseteq P_1 \boxtimes \cdots \boxtimes P_d$, implying

$$|V(T)| \leq (2r + 1)^d \leq (1 + \Delta)^0 (2r + 1)^{(0+1)(d+1)}.$$

Now assume that $k \geq 1$ and the claim holds for $k - 1$. Let \tilde{T} be the projection of $V(T)$ into H . Let (X_1, \dots, X_n) be a path decomposition of H with width $\text{pw}(H)$. We may delete any bag X_j such that $X_j \cap \tilde{T} = \emptyset$. Now assume that $X_1 \cap \tilde{T} \neq \emptyset$ and $X_n \cap \tilde{T} \neq \emptyset$. Let x be a vertex in $X_1 \cap \tilde{T}$, and let y be a vertex in $X_n \cap \tilde{T}$. Thus $(x, x_1, \dots, x_d) \in V(T)$ and $(y, y_1, \dots, y_d) \in V(T)$ for some $x_i, y_i \in V(P_i)$. Let P be the path in T with endpoints (x, x_1, \dots, x_d) and (y, y_1, \dots, y_d) . Since T has radius at most r , P has at most $2r + 1$ vertices. Let \tilde{P} be the set of vertices $v \in V(H)$ such that $(v, z_1, \dots, z_d) \in V(P)$ where $z_i \in V(P_i)$. Thus $|\tilde{P}| \leq 2r + 1$. By the choice of x and y , we have $\tilde{P} \cap X_j \neq \emptyset$ for each $j \in \{1, \dots, n\}$. Let $H' := H - \tilde{P}$. Thus $(X_1 \setminus \tilde{P}, \dots, X_n \setminus \tilde{P})$ is a path decomposition of H' with width at most $\text{pw}(H) - 1$. Let $R := \{(v, z_1, \dots, z_d) : v \in \tilde{P}, z_i \in V(P_i), i \in \{1, \dots, d\}\}$. Thus $|R| \leq (2r + 1)^{d+1}$. Let $T' := T - R$. Hence T' is a subgraph of $H' \boxtimes P_1 \boxtimes \cdots \boxtimes P_d$. Each component of T' has a neighbour in R , implying that T' has at most $\Delta |R|$ components. Every subtree of T has radius at most r (centred at the vertex closest to the centre of T). By induction, each component of T' has at most $(1 + \Delta)^{k-1} (2r + 1)^{k(d+1)}$ vertices. Thus

$$\begin{aligned} |V(T)| &\leq |R| + \Delta |R| (1 + \Delta)^{k-1} (2r + 1)^{k(d+1)} \\ &= |R| (1 + \Delta (1 + \Delta)^{k-1} (2r + 1)^{k(d+1)}) \\ &\leq |R| (1 + \Delta) (1 + \Delta)^{k-1} (2r + 1)^{k(d+1)} \\ &\leq (2r + 1)^{d+1} (1 + \Delta)^k (2r + 1)^{k(d+1)} \\ &= (1 + \Delta)^k (2r + 1)^{(k+1)(d+1)}, \end{aligned}$$

as desired. \square

Property (3) in Theorem 13 is best possible in a number of respects. First, note that we cannot allow H_1 and H_2 to have unbounded maximum degree. For example, if H_1 and H_2 are both $K_{1,n}$, then H_1 and H_2 both have pathwidth 1, but $K_{1,n} \boxtimes K_{1,n}$ contains $K_{n,n}$ as a subgraph, which contains a complete binary tree of $\Omega(\log n)$ height, which is a bounded-degree graph with exponential growth. Also, bounded pathwidth cannot be replaced by bounded treewidth, again because of the complete binary tree.

5 Polynomial Expansion

This section characterises when $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has polynomial expansion. Separators are a key tool here. A *separation* in a graph G is a pair (G_1, G_2) of subgraphs of G such that $G = G_1 \cup G_2$ and $E(G_1) \cap E(G_2) = \emptyset$. The *order* of (G_1, G_2) is $|V(G_1) \cap V(G_2)|$. A separation (G_1, G_2) is *balanced* if $|V(G_1) \setminus V(G_2)| \leq \frac{2}{3}|V(G)|$ and $|V(G_2) \setminus V(G_1)| \leq \frac{2}{3}|V(G)|$. A graph class \mathcal{G} admits *strongly sublinear separators* if there exists $c \in \mathbb{R}^+$ and $\beta \in [0, 1)$ such that for every graph $G \in \mathcal{G}$, every subgraph H of G has a balanced separation of order at most $c|V(H)|^\beta$. Dvořák and Norin [20] noted that a result of Plotkin, Rao, and Smith [44] implies that graph classes with polynomial expansion admit strongly sublinear separators. Dvořák and Norin [20] proved the converse (see [18, 22, 25] for more results on this theme).

Theorem 15 ([20]). *A hereditary class of graphs admits strongly sublinear separators if and only if it has polynomial expansion.*

Robertson and Seymour [48] established the following connection between treewidth and balanced separations.

Lemma 16 ([48, (2.6)]). *Every graph G has a balanced separation of order at most $\text{tw}(G) + 1$.*

Dvořák and Norin [23] proved the following converse.

Lemma 17 ([23]). *If every subgraph of a graph G has a balanced separation of order at most s , then $\text{tw}(G) \leq 15s$.*

We have the following strongly sublinear bound on the treewidth of graph products.

Lemma 18. *Let G be an n -vertex subgraph of $\mathbb{Z}^d \boxtimes H$ for some graph H . Then*

$$\text{tw}(G) \leq 2(\text{tw}(H) + 1)^{1/(d+1)}(dn)^{d/(d+1)} - 1.$$

Proof. Let $t := \text{tw}(H)$. For $i \in \{1, \dots, d\}$, let $\langle V_0^i, V_1^i, \dots \rangle$ be the layering of G determined by the i -th dimension. Let

$$m := \left\lceil \left(\frac{dn}{t+1} \right)^{1/(d+1)} \right\rceil.$$

For $i \in \{1, \dots, d\}$ and $\alpha \in \{0, \dots, m-1\}$, let $V^{i,\alpha} := \cup\{V_j^i : j \equiv \alpha \pmod{m}\}$. Thus $V^{i,0}, \dots, V^{i,m-1}$ is a partition of $V(G)$. Hence $|V^{i,\alpha_i}| \leq \frac{dn}{m}$ for some $\alpha_i \in \{0, \dots, m-1\}$. Let $X := \cup_{i=1}^d V^{i,\alpha_i}$. Thus $|X| \leq \frac{dn}{m}$. Note that each component of $G - X$ is a subgraph of $Q^d \boxtimes H$, where Q is the path on $m-1$ vertices. Since $\text{tw}(G)$ equals the maximum treewidth of the connected components of G , we have $\text{tw}(G) \leq \text{tw}(Q^d \boxtimes H) + |X|$. To obtain a tree decomposition of $Q^d \boxtimes H$ with width $(t+1)(m-1)^d - 1$, start with an optimal tree decomposition of H , and replace each instance of a vertex of H by the corresponding copy of Q^d . Thus

$$\text{tw}(G) \leq (t+1)(m-1)^d - 1 + \frac{dn}{m} \leq 2(t+1)^{1/(d+1)}(dn)^{d/(d+1)} - 1. \quad \square$$

Lemma 18 is generalised by our next result, which characterises when a graph product has polynomial expansion. The following definition is key. Say that graph classes \mathcal{G}_1 and \mathcal{G}_2 have *joint polynomial growth* if there exists a polynomial function p such that for every $r \in \mathbb{N}$, there exists $i \in \{1, 2\}$ such that for every graph $G \in \mathcal{G}_i$ every r -ball in G has size at most $p(r)$.

Theorem 19. *The following are equivalent for hereditary graph classes \mathcal{G}_1 and \mathcal{G}_2 :*

- (1) $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has polynomial expansion,
- (2) $\mathcal{G}_1 \square \mathcal{G}_2$ has polynomial expansion,
- (3) \mathcal{G}_1 has polynomial expansion, \mathcal{G}_2 has polynomial expansion, and \mathcal{G}_1 and \mathcal{G}_2 have joint polynomial growth.

Proof. (1) implies (2) since $\mathcal{G}_1 \square \mathcal{G}_2 \subseteq \mathcal{G}_1 \boxtimes \mathcal{G}_2$.

We now show that (2) implies (3). Assume that $\mathcal{G}_1 \square \mathcal{G}_2$ has polynomial expansion. That is, for some polynomial g , for every graph $G \in \mathcal{G}_1 \square \mathcal{G}_2$, every r -shallow minor of G has average degree at most $g(r)$. Since $\mathcal{G}_1 \cup \mathcal{G}_2 \subseteq \mathcal{G}_1 \square \mathcal{G}_2$, both \mathcal{G}_1 and \mathcal{G}_2 have polynomial expansion.

Assume for the sake of contradiction that \mathcal{G}_1 and \mathcal{G}_2 do not have joint polynomial growth. Thus for every polynomial p there exists $r \in \mathbb{N}$ such that for each $i \in \{1, 2\}$ some r -ball of some graph $G_i \in \mathcal{G}_i$ has at least $p(r)$ vertices. Apply this where p is a polynomial with $p(r) \geq \max\{1 + rn, \binom{n}{2}\}$, where $n := \lceil g(2r) + 2 \rceil$. Since \mathcal{G}_1 and \mathcal{G}_2 are hereditary, there exists $r \in \mathbb{N}$ such that there is a graph $G_1 \in \mathcal{G}_1$ with radius at most r and at least $1 + rn$ vertices, and there is a graph $G_2 \in \mathcal{G}_2$ with radius at most r and at least $\binom{n}{2}$ vertices.

Let z be the central vertex in G_1 . Since $|V(G_1)| \geq 1 + rn$, for some $i \in \{1, \dots, r\}$, there is a set A of n vertices in G_1 at distance exactly i from z . For all $\{v, w\} \in \binom{A}{2}$, let $P_{v,w}$ be the shortest vw -path contained with the union of a shortest vz -path and a shortest wz -path in G_1 . Thus $P_{v,w}$ has length at most $2r$ and $V(P_{v,w}) \cap A = \{v, w\}$. Let B be a set of $\binom{n}{2}$ vertices in G_2 . Fix an arbitrary bijection $\sigma : \binom{A}{2} \rightarrow B$.

Let $G := G_1 \square G_2$. For each $v \in A$, let $X_v := G[\{(v, x) : x \in V(G_2)\}]$; note that X_v is isomorphic to G_2 , and thus has radius at most r . Moreover, X_v and X_w are disjoint for distinct $v, w \in A$. For $\{v, w\} \in \binom{A}{2}$, let $Y_{v,w} := G[\{(x, \sigma(\{v, w\})) : x \in V(G_1)\}]$; note that $Y_{v,w}$ is isomorphic to G_1 . Let $Q_{v,w}$ be the copy of the path $P_{v,w}$ within $Y_{v,w}$. Since $V(P_{v,w}) \cap A = \{v, w\}$, the only vertices of $Q_{v,w}$ in $\bigcup_{u \in A} X_u$ are $(v, \sigma(\{v, w\}))$ and $(w, \sigma(\{v, w\}))$, which are the endpoints of $Q_{v,w}$ in X_v and X_w respectively. Since $P_{v,w}$ has length at most $2r$, so does $Q_{v,w}$.

By construction, $Q_{v,w}$ and $Q_{p,q}$ are disjoint for distinct $\{v, w\}, \{p, q\} \in \binom{A}{2}$. Contract X_v to a vertex for each $v \in A$, and contract $Q_{v,w}$ to an edge for each $\{v, w\} \in \binom{A}{2}$. We obtain the complete graph K_n as a minor in G . Moreover, the minor is $2r$ -shallow. This is a contradiction, since K_n has average degree greater than $g(2r)$.

We prove that (3) implies (1) by a series of lemmas below (culminating in Lemma 23 below). \square

For a graph G , a set $X \subseteq V(G)$ is r -localising if for every component C of $G - X$, there exists a vertex $v \in V(G)$ such that $d_G(u, v) < r$ for every $u \in C$ (note that the distance is in G , not in $G - X$).

The following is a variation on Lemma 5.2 of [36]. For $r \in \mathbb{N}$ and $p, q \in \mathbb{R}$ with $0 < p, q < 1$, consider the following function $f_{r,p,q}$ defined on $\{0, 1, \dots, r\}$. First, let $f_{r,p,q}(r) := p$. Now, for every integer $s \in \{0, 1, \dots, r-1\}$, inductively define

$$f_{r,p,q}(s) := \min(q f_{r,p,q}(\{s+1, \dots, r\}), 1 - f_{r,p,q}(\{s+1, \dots, r\})),$$

where $f(S) := \sum_{i \in S} f(i)$.

Lemma 20. Fix $r \in \mathbb{N}$ and $p, q \in \mathbb{R}$ with $0 < p, q < 1$, such that $f_{r,p,q}(\{0, 1, \dots, r\}) = 1$ (so $f_{r,p,q}$ defines a probability distribution on $\{0, 1, \dots, r\}$). For every graph G , there exists a probability distribution over the r -localising subsets of $V(G)$ such that the set X drawn from this distribution satisfies $\mathbb{P}[v \in X] \leq p|N^r(v)| + q$ for every $v \in V(G)$.

Proof. Let $V(G) = \{v_1, \dots, v_n\}$. For $i \in \{1, \dots, n\}$, choose $r_i \in \{0, 1, \dots, r\}$ independently at random such that $\mathbb{P}[r_i = s] = f_{r,p,q}(s)$. For each $x \in V(G)$, let $i(x)$ be the minimum index i such that $d(x, v_i) \leq r_i$, and let $X := \{x \in V(G) : d(x, v_{i(x)}) = r_{i(x)}\}$.

First we argue that X is r -localising. Consider any component C of $G - X$, and let z be the vertex of C with $i(z)$ minimum. Suppose for the sake of contradiction that C contains a vertex u at distance at least r from $v_{i(z)}$, and let P be a path from z to u in C . Then P contains a vertex x at distance exactly $r_{i(z)}$ from $v_{i(z)}$. However, since $i(x) \geq i(z)$, we conclude $i(x) = i(z)$ and $x \in X$, which is a contradiction.

Next, we bound the probability that a vertex v of G is in X . Consider any $i \in \{1, \dots, n\}$. If $d(v, v_i) > r$, then $\mathbb{P}[i(v) = i] = 0$. If $d(v, v_i) = r$, then $\mathbb{P}[i(v) = i] \leq \mathbb{P}[r_{i(v)} = r] = p$. If $d(v, v_i) < r$, then letting $s := d(v, v_i)$ we have

$$\begin{aligned} \mathbb{P}[v \in X | i(v) = i] &= \mathbb{P}[r_i = s | r_1 < d(v, v_1), \dots, r_{i-1} < d(v, v_{i-1}), r_i \geq s] \\ &= \mathbb{P}[r_i = s | r_i \geq s] \\ &= \frac{f_{r,p,q}(s)}{f_{r,p,q}(\{s, \dots, r\})} \\ &\leq q. \end{aligned}$$

Therefore,

$$\mathbb{P}[v \in X] = \sum_{i=1}^n \mathbb{P}[i(v) = i] \cdot \mathbb{P}[v \in X | i(v) = i] \leq p|N^r(v)| + q. \quad \square$$

Corollary 21. For every polynomial g , there exists r_0 such that the following holds. Let $r \geq r_0$ be a positive integer and let G be a graph such that $|N^r(v)| \leq g(r)$ for every $v \in V(G)$. Then there exists a probability distribution over the r -localising subsets of $V(G)$ such that the set X drawn from this distribution satisfies $\mathbb{P}[v \in X] \leq 2r^{-1/2}$ for every $v \in V(G)$.

Proof. Let c be the degree of g plus one, so that $g(r) \leq r^c$ for every sufficiently large r . Let $p := r^{-c-1/2}$ and $q := r^{-1/2}$. Note that for sufficiently large r ,

$$p(1+q)^r \geq pe^{qr/2} = \exp(\sqrt{r}/2 - (c+1/2)\log r) > 1.$$

Hence $f_{r,p,q}(r) = p > 1/(q+1)^r$. It follows by induction that $f_{r,p,q}(\{s, \dots, r\}) \geq 1/(q+1)^s$ for each $s \in \{1, \dots, r\}$. Thus $f_{r,p,q}(0) = 1 - f_{r,p,q}(\{1, \dots, r\})$ and $f_{r,p,q}(\{0, \dots, r\}) = 1$. The claim follows from Lemma 20. \square

Corollary 22. *For every polynomial g , there exists r_0 such that the following holds. Let $r \geq r_0$ be a positive integer and let G be a graph such that $|N^r[v]| \leq g(r)$ for every $v \in V(G)$. Then for every function $w : V(G) \rightarrow \mathbb{R}_0^+$, there exists $X \subseteq V(G)$ such that $w(X) \leq 2r^{-1/2}w(V(G))$ and each component of $G - X$ has at most $g(r)$ vertices.*

Proof. Choose an r -localising set $X \subseteq V(G)$ using Corollary 21. Since X is r -localising and $|N^r[v]| \leq g(r)$ for every $v \in V(G)$, each component of $G - X$ has at most $g(r)$ vertices. Furthermore,

$$\mathbb{E}[|w(X)|] = \sum_{v \in V(G)} \mathbb{P}[v \in X]w(v) \leq 2r^{-1/2}w(V(G)).$$

Hence there is a choice for X such that $w(X) \leq 2r^{-1/2}w(V(G))$. \square

Lemma 23. *Suppose \mathcal{G}_1 and \mathcal{G}_2 are classes with strongly sublinear separators and of joint polynomial growth (bounded by a polynomial g). Then $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has strongly sublinear separators.*

Proof. Let $\varepsilon > 0$ be such that every subgraph F of a graph from $\mathcal{G}_1 \cup \mathcal{G}_2$ has a balanced separator of order at most $\lceil |V(F)|^{1-\varepsilon} \rceil$. Let $\beta > 0$ be sufficiently small (depending on ε and g).

Suppose $G_1 \in \mathcal{G}_1$, $G_2 \in \mathcal{G}_2$, and H is a subgraph of $G_1 \boxtimes G_2$. Let π_1 and π_2 be the projections from H to G_1 and G_2 . Let $n := |V(H)|$ and $r := n^\beta$. By symmetry, we may assume $|N^r[v]| \leq g(r)$ for every vertex v of G_1 . Let $w(v) := |\pi_1^{-1}(v)|$ for each $v \in V(G_1)$. By Corollary 22, there exists $X \subseteq V(G_1)$ such that $w(X) = O(r^{-1/2}n) = O(n^{1-\beta/2})$ and each component of $G_1 - X$ has at most $g(r) = g(n^\beta) = O(n^{\varepsilon/2})$ vertices. Let $A := \pi_1^{-1}(X)$; then $|A| = w(X) = O(n^{1-\beta/2})$.

The graph G_2 has treewidth $O(n^{1-\varepsilon})$, and thus the product of G_2 with $G_1 - X$ (as well as its subgraph $H - A$) has treewidth $O(n^{1-\varepsilon}g(r)) = O(n^{1-\varepsilon/2})$. Consequently, $H - A$ has a balanced separator B of order $O(n^{1-\varepsilon/2})$, and $A \cup B$ is a balanced separator of H of order $O(n^{1-\min(\varepsilon, \beta)/2})$. \square

6 Bounded Expansion

This section characterises when $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has bounded expansion. The following definition by Kierstead and Yang [35] is the key tool. For a graph G , linear ordering \preceq of $V(G)$, vertex $v \in V(G)$, and integer $r \geq 1$, a vertex x is (r, \preceq) -reachable from v if there is a path $v = v_0, v_1, \dots, v_{r'} = x$ of length $r' \in \{0, 1, \dots, r\}$ such that $x \preceq v \prec v_i$ for all $i \in \{1, 2, \dots, r' - 1\}$. For a graph G and $r \in \mathbb{N}$, the r -colouring number $\text{col}_r(G)$ of G , also known as the *strong r -colouring number*, is the minimum integer k such that there is a linear ordering \preceq of $V(G)$ such that at most k vertices are (r, \preceq) -reachable from each vertex v of G . For example, van den Heuvel, Ossona de Mendez, Quiroz, Rabinovich, and Siebertz [50] proved that every planar graph G satisfies $\text{col}_r(G) \leq 5r + 1$, and more generally, that every K_t -minor-free graph G satisfies $\text{col}_r(G) \leq \binom{t-1}{2}(2r + 1)$. Most generally, Zhu [54] showed that these r -colouring numbers characterise bounded expansion classes.

Theorem 24 ([54]). *A graph class \mathcal{G} has bounded expansion if and only if for each $r \in \mathbb{N}$ there exists $c \in \mathbb{N}$ such that $\text{col}_r(G) \leq c$ for all $G \in \mathcal{G}$.*

We now show that if G has bounded r -colouring number and H has bounded maximum degree, then $G \boxtimes H$ has bounded r -colouring number.

Lemma 25. *If G is a graph with $\text{col}_r(G) \leq c$ and H is a graph with maximum degree at most Δ , then $\text{col}_r(G \boxtimes H) < c(\Delta + 2)^r$.*

Proof. Let G^+ and H^+ be the pseudographs obtained from G and H by adding a loop at every vertex. Let \preceq_G be a vertex-ordering of G witnessing that $\text{col}_r(G) \leq c$. Let \preceq be an ordering of $V(G \boxtimes H)$ where $v_1 \prec_G v_2$ implies $(v_1, w_1) \prec (v_2, w_2)$ for all $v_1, v_2 \in V(G)$ and $w_1, w_2 \in V(H)$. We now bound the number of vertices of $G \boxtimes H$ that are (r, \preceq) -reachable from a fixed vertex $(v, w) \in V(G \boxtimes H)$. Say (x, y) is (r, \preceq) -reachable from (v, w) . Thus there is a path $(v, w) = (v_0, w_0), (v_1, w_1), \dots, (v_{r'}, w_{r'}) = (x, y)$ of length $r' \in \{0, \dots, r\}$, such that $(v_{r'}, w_{r'}) \preceq (v, w) \prec (v_i, w_i)$ for each $i \in \{1, \dots, r' - 1\}$. Charge (x, y) to the pair $(x, (w_0, w_1, \dots, w_{r'}))$. By the definition of \boxtimes , the sequence $(v_0, v_1, \dots, v_{r'})$ is a walk in G^+ , and the sequence $(w_0, w_1, \dots, w_{r'})$ is a walk in H^+ . By the definition of \preceq , we have $v_{r'} \preceq v_0 \preceq v_i$ for each $i \in \{1, \dots, r' - 1\}$. Thus $v_{r'}$ is (r, \preceq_G) -reachable from v_0 in G . By assumption, at most c vertices are (r, \preceq_G) -reachable from v_0 in G . The number of walks of length at most r in H^+ starting at w_0 is at most $\sum_{i=0}^r (\Delta + 1)^i < (\Delta + 2)^r$. Thus for each vertex $x \in V(G)$, less than $(\Delta + 2)^r$ vertices (r, \preceq) -reachable from (v, w) are charged to some pair (x, W) . Hence, less than $c(\Delta + 2)^r$ vertices in $G \boxtimes H$ are (r, \preceq_G) -reachable from (v, w) in \preceq . Therefore $\text{col}_r(G \boxtimes H) < c(\Delta + 2)^r$. \square

The next theorem is the main contribution of this section.

Theorem 26. *The following are equivalent for hereditary graph classes \mathcal{G}_1 and \mathcal{G}_2 :*

1. $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has bounded expansion,
2. $\mathcal{G}_1 \square \mathcal{G}_2$ has bounded expansion,
3. both \mathcal{G}_1 and \mathcal{G}_2 have bounded expansion, and at least one of \mathcal{G}_1 and \mathcal{G}_2 has bounded maximum degree.

Proof. (1) implies (2) since $G_1 \square G_2 \subseteq G_1 \boxtimes G_2$.

We now show that (2) implies (3). Assume that $\mathcal{G}_1 \square \mathcal{G}_2$ has bounded expansion. Since $\mathcal{G}_1 \cup \mathcal{G}_2 \subseteq \mathcal{G}_1 \boxtimes \mathcal{G}_2$, both \mathcal{G}_1 and \mathcal{G}_2 also have bounded expansion. If neither \mathcal{G}_1 nor \mathcal{G}_2 have bounded maximum degree, then for every $n \in \mathbb{N}$, the star graph $K_{1,n}$ is a subgraph of some graph $G_1 \in \mathcal{G}_1$ and of some graph $G_2 \in \mathcal{G}_2$. Observe that $K_{1,n} \square K_{1,n}$ contains a 1-subdivision of $K_{n,n}$. Thus $G_1 \square G_2$ contains a graph with average degree n (namely, $K_{n,n}$) as a 1-shallow minor, which is a contradiction since $\mathcal{G}_1 \square \mathcal{G}_2$ has bounded expansion. Hence at least one of \mathcal{G}_1 and \mathcal{G}_2 has bounded maximum degree.

We now show that (3) implies (1). Assume that both \mathcal{G}_1 and \mathcal{G}_2 have bounded expansion, and every graph in \mathcal{G}_2 has maximum degree at most Δ . By Theorem 24, for each $r \in \mathbb{N}$ there exists $c_r \in \mathbb{N}$ such that $\text{col}_r(G) \leq c_r$ for all $G_1 \in \mathcal{G}_1$. Let $G_2 \in \mathcal{G}_2$. By Lemma 25, we have $\text{col}_r(G_1 \boxtimes G_2) \leq c_r(\Delta + 2)^r$, and the same bound holds for every subgraph of $G_1 \boxtimes G_2$. By Theorem 24, $\mathcal{G}_1 \boxtimes \mathcal{G}_2$ has bounded expansion. \square

7 Geometric Graph Classes

The section explores graph product structure theorems for various geometrically defined graph classes.

The *unit disc* graph of a finite set $X \subseteq \mathbb{R}^d$ is the graph G with $V(G) = X$ where $vw \in E(G)$ if and only if $\text{dist}(v, w) \leq 1$. Here dist is the Euclidean distance in \mathbb{R}^d . Let \mathbb{Z}^d be the strong product of d paths $P \boxtimes \dots \boxtimes P$ (the d -dimensional grid graph with crosses). The next result describes unit discs in terms of strong products, which implies that the class of unit disc graphs with bounded dimension and bounded maximum clique size has polynomial growth.

Theorem 27. *Every unit disc graph G in \mathbb{R}^d with no $(k+1)$ -clique is a subgraph of $\mathbb{Z}^d \boxtimes K_{k\lceil\sqrt{d}\rceil^d}$.*

Proof. Let $t := k\lceil\sqrt{d}\rceil^d$. Let $x_i(v)$ be the i -th coordinate of each vertex $v \in V(G)$. For $p_1, \dots, p_d \in \mathbb{Z}$, let $V\langle p_1, \dots, p_d \rangle$ be the set of vertices $v \in V(G)$ such that $p_i \leq x_i(v) < p_i + 1$ for each $i \in \{1, \dots, d\}$. Thus the sets $V\langle p_1, \dots, p_d \rangle$ partition $V(G)$. Each set $V\langle p_1, \dots, p_d \rangle$ consists of the set of vertices in a particular unit cube. Note that the unit cube can be partitioned into $\lceil\sqrt{d}\rceil^d$ sub-cubes, each with side length at most $\frac{1}{\sqrt{d}}$ and thus with diameter at most 1. The set of vertices in a sub-cube with diameter at most 1 is a clique in G . Thus at most k vertices lie in a single sub-cube, and $|V\langle p_1, \dots, p_d \rangle| \leq t$. Injectively label the vertices in $V\langle p_1, \dots, p_d \rangle$ by $1, 2, \dots, t$. Map each vertex v in $V\langle p_1, \dots, p_d \rangle$ labelled $\ell(v)$ to the vertex $(p_1, \dots, p_d, \ell(v))$ of $\mathbb{Z}^d \boxtimes K_t$. Thus the vertices of G are mapped to distinct vertices of $\mathbb{Z}^d \boxtimes K_t$. For each edge $vw \in E(G)$, if $v \in V\langle p_1, \dots, p_d \rangle$ and $w \in V\langle q_1, \dots, q_d \rangle$, then $|p_i - q_i| \leq 1$ for each $i \in \{1, \dots, d\}$, and if $p_i = q_i$ for each $i \in \{1, \dots, d\}$, then $\ell(v) \neq \ell(w)$. Thus v and w are mapped to adjacent vertices in $\mathbb{Z}^d \boxtimes K_t$. \square

By a volume argument, every covering of the unit cube by balls of diameter 1 uses at least $(\frac{d}{18})^{d/2}$ balls. So the $k\lceil\sqrt{d}\rceil^d$ term in the above theorem cannot be drastically improved.

The *k-nearest neighbour graph* of a finite set $P \subset \mathbb{R}^d$ has vertex set P , where two vertices v and w are adjacent if w is the one of the k points in P closest to v , or v is the one of the k points in P closest to w . Miller, Teng, Thurston, and Vavasis [38] showed that such graphs admit separators of order $O(n^{1-1/d})$.

Can we describe the structure of k -nearest-neighbour graphs using graph products?

Conjecture 28. Every k -nearest neighbour graph in \mathbb{R}^d is a subgraph of $H \boxtimes \mathbb{Z}^{d-1}$ for some graph H with treewidth at most $f(k, d)$.

This conjecture is trivial for $d = 1$ and true for $d = 2$, as proved by Dujmović et al. [17]. Note that “treewidth at most $f(k, d)$ ” cannot be replaced by “pathwidth at most $f(k, d)$ ” for $d \geq 2$ because complete binary trees are 2-dimensional 2-nearest neighbour graphs without polynomial growth (see Theorem 13).

Here is a still more general example: Miller et al. [38] defined a (d, k) -neighbourhood system to consist of a collection \mathcal{C} of n balls in \mathbb{R}^d such that no point in \mathbb{R}^d is covered by more than k balls. Consider the associated graph with one vertex for each ball, where two vertices are adjacent if the corresponding balls intersect. Miller et al. [38] showed that such graphs admit balanced separators of order $O(n^{1-1/d})$. Note that by the Koebe circle packing theorem, every planar graph is associated with some $(2, 2)$ -neighbourhood system. Thus the result of Miller et al. [38] is a far-reaching generalisation of the Lipton-Tarjan Separator Theorem [37]. Is there a product structure theorem for these graphs? Might the structure in Theorem 4 be applicable here?

Open Problem 29. If G is the graph associated with a (d, k) -neighbourhood system, can G be obtained from clique-sums of graphs G_1, \dots, G_n such that $G_i \subseteq (H_i \boxtimes P^{(d-1)}) + K_a$, for some graph H_i with treewidth at most k , where a is a constant that depends only on k and d . The natural choice is $a = k - 1$.

One can ask a similar question for graphs embeddable in a finite-dimensional Euclidean space with bounded distortion of distances. Dvořák [22] showed that such graphs have strongly sublinear separators.

8 Open Problems

We finish the paper with a number of open problems.

It is open whether the treewidth 4 bound in Theorem 2(b) can be improved.

Open Problem 30. For every $g \in \mathbb{N}$, does there exist $t \in \mathbb{N}$ such that every graph of Euler genus g is a subgraph of $H \boxtimes P \boxtimes K_t$ for some graph H of treewidth at most 3?

The proofs of Theorems 3 and 4 both use the Graph Minor Structure Theorem of Robertson and Seymour [49].

Open Problem 31. *Is there a proof of Theorem 3 or Theorem 4 that does not use the graph minor structure theorem?*

The following problem asks to minimise the treewidth in Theorem 7.

Open Problem 32 ([17]). *Does there exist a constant c such that for every $k \in \mathbb{N}$ there exists $t \in \mathbb{N}$ such that every k -planar graph is a subgraph of $H \boxtimes P \boxtimes K_t$ for some graph H of treewidth at most c ?*

Open Problem 33. *Can any graph class with linear or polynomial expansion be described as a product of simpler graph classes along with apex vertices, clique sums, and other ingredients.*

Such a theorem would be useful for proving properties about such classes. Recent results say that the “other ingredients” in Open Problem 33 are needed, as we now explain. Let G' be the $6\text{tw}(G)$ -subdivision of a graph G . Let $\mathcal{G} := \{G' : G \text{ is a graph}\}$. Grohe, Kreutzer, Rabinovich, Siebertz, and Stavropoulos [30] proved that \mathcal{G} has linear expansion. On the other hand, Dubois, Joret, Perarnau, Pilipczuk, and Pitois [11] proved that there are graphs G such that every p -centred colouring of G' has at least $2^{cp^{1/2}}$ colours, for some constant $c > 0$. We do not define “ p -centred colouring” here since we do not need the definition. All we need to know is that subgraphs of $H \boxtimes P$, where H has bounded treewidth and P is a path, have p -centred colourings with $f(p)$ colours, where f is a polynomial function (see [10, 17]). This result is easily extended to allow for apex vertices and clique sums. This shows that there are graphs with linear expansion that cannot be described solely in terms of products of bounded treewidth graphs and paths (plus apex vertices and clique sums). For related results, see [19].

One way to test the quality of such a structure theorem is whether they resolve the following questions about queue-number and nonrepetitive chromatic number mentioned in Section 1:

Open Problem 34. *Do graph classes with linear or polynomial expansion have bounded queue-number?*

Open Problem 35. *Do graphs classes with linear or polynomial (or even single exponential) expansion have bounded nonrepetitive chromatic number?*

Note that bounded degree graphs are an example with exponential expansion and unbounded queue-number [51]. Similarly, subdivisions of complete graphs K_n with $o(\log n)$ division vertices per edge are an example with super-exponential expansion and unbounded nonrepetitive chromatic number [41]. Thus the graph classes mentioned in the above open problems are the largest possible with bounded queue-number or bounded nonrepetitive chromatic number.

8.1 Algorithmic Questions

Do product structure theorems have algorithmic applications? Consider the method of Baker [2] for designing polynomial-time approximation schemes for problems on planar graphs. This method partitions the graph into BFS layers, such that the problem can be solved optimally on each layer (since the induced subgraph has bounded treewidth), and then combines the solutions from each layer. Theorem 1 gives a more precise description of the layered structure of planar graphs. It is conceivable that this extra structural information is useful when designing algorithms for planar graphs (and any graph class that has a product structure theorem).

Some NP-complete problems can be solved efficiently on planar graphs. Can these results be extended to any subgraph of the strong product of a bounded treewidth graph and a path? For example, can max-cut be solved efficiently for graphs that are subgraphs of $H \boxtimes P$, where H is a bounded treewidth graph and P is a path, such as apex-minor-free graphs? This would be a considerable generalisation of the known polynomial-time algorithm for max-cut on planar graphs [31] and on graphs of bounded genus [29].

Some problems can be solved by particularly fast algorithms on planar graphs. Can such results be generalised for any subgraph of the strong product of a bounded treewidth graph and a path? For example, can shortest paths be computed in $O(n)$ time for n -vertex subgraphs of $H \boxtimes P$, where H is a bounded treewidth graph and P is a path? Can maximum flows be computed in $n \log^{O(1)}(n)$ time for n -vertex subgraphs of $H \boxtimes P$? See [6, 24] for analogous results for planar graphs.

References

- [1] NOGA ALON, JAROSŁAW GRYTCZUK, MARIUSZ HAŁUSZCZAK, AND OLIVER RIORDAN. Nonrepetitive colorings of graphs. *Random Structures Algorithms*, 21(3–4):336–346, 2002. doi: 10.1002/rsa.10057. MR: 1945373.
- [2] BRENDA S. BAKER. Approximation algorithms for NP-complete problems on planar graphs. *J. ACM*, 41(1):153–180, 1994. doi: 10.1145/174644.174650. MR: 1369197.
- [3] HANS L. BODLAENDER. A tourist guide through treewidth. *Acta Cybernet.*, 11(1-2):1–21, 1993.
- [4] HANS L. BODLAENDER. A partial k -arboretum of graphs with bounded treewidth. *Theoret. Comput. Sci.*, 209(1-2):1–45, 1998. doi: 10.1016/S0304-3975(97)00228-4. MR: 1647486.
- [5] MARTHE BONAMY, CYRIL GAVOILLE, AND MICHAŁ PILIPCZUK. Shorter labeling schemes for planar graphs. In SHUCHI CHAWLA, ed., *Proc. ACM-SIAM Symposium on Discrete Algorithms (SODA '20)*, pp. 446–462. 2020. doi: 10.1137/1.9781611975994.27. arXiv: 1908.03341.

- [6] SERGIO CABELLO, ERIN W. CHAMBERS, AND JEFF ERICKSON. Multiple-source shortest paths in embedded graphs. *SIAM J. Comput.*, 42(4):1542–1571, 2013. doi: 10.1137/120864271.
- [7] ZHI-ZHONG CHEN, MICHELANGELO GRIGNI, AND CHRISTOS H. PAPADIMITRIOU. Map graphs. *J. ACM*, 49(2):127–138, 2002. doi: 10.1145/506147.506148. MR: 2147819.
- [8] REINHARD DIESTEL. *Graph theory*, vol. 173 of *Graduate Texts in Mathematics*. Springer, 4th edn., 2010. <http://diestel-graph-theory.com/>. MR: 2744811.
- [9] GUOLI DING AND BOGDAN OPOROWSKI. Some results on tree decomposition of graphs. *J. Graph Theory*, 20(4):481–499, 1995. doi: 10.1002/jgt.3190200412. MR: 1358539.
- [10] MICHAŁ DĘBSKI, STEFAN FELSNER, PIOTR MICEK, AND FELIX SCHRÖDER. Improved bounds for centered colorings. In SHUCHI CHAWLA, ed., *Proc. ACM-SIAM Symposium on Discrete Algorithms (SODA '20)*, pp. 2212–2226. 2020. doi: 10.1137/1.9781611975994.136. arXiv: 1907.04586.
- [11] LOIČ DUBOIS, GWENAËL JORET, GUILLEM PERARNAU, MARCIN PILIPCZUK, AND FRANÇOIS PITOIS. Two lower bounds for p -centered colorings. 2020. arXiv: 2006.04113.
- [12] VIDA DUJMOVIĆ, DAVID EPPSTEIN, GWENAËL JORET, PAT MORIN, AND DAVID R. WOOD. Minor-closed graph classes with bounded layered pathwidth. *SIAM J. Disc. Math.*, to appear. arXiv: 1810.08314.
- [13] VIDA DUJMOVIĆ, DAVID EPPSTEIN, AND DAVID R. WOOD. Structure of graphs with locally restricted crossings. *SIAM J. Discrete Math.*, 31(2):805–824, 2017. doi: 10.1137/16M1062879.
- [14] VIDA DUJMOVIĆ, LOUIS ESPERET, PAT MORIN, BARTOSZ WALCZAK, AND DAVID R. WOOD. Clustered 3-colouring graphs of bounded degree. 2020. arXiv: 2002.11721.
- [15] VIDA DUJMOVIĆ, GWENAËL JORET, PIOTR MICEK, PAT MORIN, TORSTEN UECKERDT, AND DAVID R. WOOD. Planar graphs have bounded queue-number. In *Proc. 60th Annual Symp. Foundations Comput. Sci. (FOCS '19)*, pp. 862–875. IEEE, 2019. doi: 10.1109/FOCS.2019.00056.
- [16] VIDA DUJMOVIĆ, GWENAËL JORET, PIOTR MICEK, PAT MORIN, TORSTEN UECKERDT, AND DAVID R. WOOD. Planar graphs have bounded queue-number. *J. ACM*, 2020. doi: 10.1145/3385731.
- [17] VIDA DUJMOVIĆ, PAT MORIN, AND DAVID R. WOOD. Graph product structure for non-minor-closed classes, 2019. arXiv: 1907.05168.
- [18] ZDENĚK DVOŘÁK. Sublinear separators, fragility and subexponential expansion. *European J. Combin.*, 52(A):103–119, 2016. doi: 10.1016/j.ejc.2015.09.001.
- [19] ZDENĚK DVOŘÁK, ROSE MCCARTY, AND SERGEY NORIN. Sublinear separators in intersection graphs of convex shapes. 2020. arXiv: 2001.01552.
- [20] ZDENĚK DVOŘÁK AND SERGEY NORIN. Strongly sublinear separators and polynomial expansion. *SIAM J. Discrete Math.*, 30(2):1095–1101, 2016. doi: 10.1137/15M1017569. MR: 3504982.

- [21] ZDENĚK DVOŘÁK AND ROBIN THOMAS. List-coloring apex-minor-free graphs. 2014. arXiv: 1401.1399.
- [22] ZDENĚK DVOŘÁK. On classes of graphs with strongly sublinear separators. *European J. Combin.*, 71:1–11, 2018. doi: 10.1016/j.ejc.2018.02.032. MR: 3802229.
- [23] ZDENĚK DVOŘÁK AND SERGEY NORIN. Treewidth of graphs with balanced separations. *J. Combin. Theory Ser. B*, 137:137–144, 2019. doi: 10.1016/j.jctb.2018.12.007. MR: 3980088.
- [24] JEFF ERICKSON. Maximum flows and parametric shortest paths in planar graphs. In MOSES CHARIKAR, ed., *Proc. 21st Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2010*, pp. 794–804. SIAM, 2010. doi: 10.1137/1.9781611973075.65.
- [25] LOUIS ESPERET AND JEAN-FLORENT RAYMOND. Polynomial expansion and sublinear separators. *European J. Combin.*, 69:49–53, 2018. doi: 10.1016/j.ejc.2017.09.003.
- [26] FEDOR V. FOMIN, DANIEL LOKSHTANOV, AND SAKET SAURABH. Bidimensionality and geometric graphs. In *Proc. 23rd Annual ACM-SIAM Symposium on Discrete Algorithms*, pp. 1563–1575. 2012. doi: 10.1137/1.9781611973099.124. MR: 3205314.
- [27] JACOB FOX AND JÁNOS PACH. A separator theorem for string graphs and its applications. *Combin. Probab. Comput.*, 19(3):371–390, 2010. doi: 10.1017/S0963548309990459.
- [28] JACOB FOX AND JÁNOS PACH. Applications of a new separator theorem for string graphs. *Combin. Probab. Comput.*, 23(1):66–74, 2014. doi: 10.1017/S0963548313000412.
- [29] ANNA GALLUCCIO AND MARTIN LOEBL. On the theory of Pfaffian orientations. II. T -joins, k -cuts, and duality of enumeration. *Electron. J. Combin.*, 6(1):#R7, 1999. doi: 10.37236/1439.
- [30] MARTIN GROHE, STEPHAN KREUTZER, ROMAN RABINOVICH, SEBASTIAN SIEBERTZ, AND KONSTANTINOS STAVROPOULOS. Coloring and covering nowhere dense graphs. *SIAM J. Discrete Math.*, 32(4):2467–2481, 2018. doi: 10.1137/18M1168753.
- [31] FRANK HADLOCK. Finding a maximum cut of a planar graph in polynomial time. *SIAM J. Comput.*, 4(3):221–225, 1975. doi: 10.1137/0204019.
- [32] DANIEL J. HARVEY AND DAVID R. WOOD. Parameters tied to treewidth. *J. Graph Theory*, 84(4):364–385, 2017. doi: 10.1002/jgt.22030. MR: 3623383.
- [33] LENWOOD S. HEATH, F. THOMSON LEIGHTON, AND ARNOLD L. ROSENBERG. Comparing queues and stacks as mechanisms for laying out graphs. *SIAM J. Discrete Math.*, 5(3):398–412, 1992. doi: 10.1137/0405031. MR: 1172748.
- [34] SAMPATH KANNAN, MONI NAOR, AND STEVEN RUDICH. Implicit representation of graphs. *SIAM J. Discrete Math.*, 5(4):596–603, 1992. doi: 10.1137/0405049.

- [35] HAL A. KIERSTEAD AND DAQING YANG. Orderings on graphs and game coloring number. *Order*, 20(3):255–264, 2003. doi: 10.1023/B:ORDE.0000026489.93166.cb.
- [36] ROBERT KRAUTHGAMER AND JAMES R. LEE. The intrinsic dimensionality of graphs. *Combinatorica*, 27(5):551–585, 2007. doi: 10.1007/s00493-007-2183-y.
- [37] RICHARD J. LIPTON AND ROBERT E. TARJAN. A separator theorem for planar graphs. *SIAM J. Appl. Math.*, 36(2):177–189, 1979. doi: 10.1137/0136016. MR: 0524495.
- [38] GARY L. MILLER, SHANG-HUA TENG, WILLIAM THURSTON, AND STEPHEN A. VAVASIS. Separators for sphere-packings and nearest neighbor graphs. *J. ACM*, 44(1):1–29, 1997. doi: 10.1145/256292.256294. MR: 1438463.
- [39] BOJAN MOHAR AND CARSTEN THOMASSEN. *Graphs on surfaces*. Johns Hopkins University Press, 2001. MR: 1844449.
- [40] JAROSLAV NEŠETŘIL AND PATRICE OSSONA DE MENDEZ. *Sparsity*, vol. 28 of *Algorithms and Combinatorics*. Springer, 2012. doi: 10.1007/978-3-642-27875-4. MR: 2920058.
- [41] JAROSLAV NEŠETŘIL, PATRICE OSSONA DE MENDEZ, AND DAVID R. WOOD. Characterisations and examples of graph classes with bounded expansion. *European J. Combin.*, 33(3):350–373, 2011. doi: 10.1016/j.ejc.2011.09.008. MR: 2864421.
- [42] JÁNOS PACH AND GÉZA TÓTH. Recognizing string graphs is decidable. *Discrete Comput. Geom.*, 28(4):593–606, 2002. doi: 10.1007/s00454-002-2891-4.
- [43] TARIK PECANINOVIC. Complete graph minors in strong products. 2019. Honours thesis, School of Mathematics, Monash University.
- [44] SERGE PLOTKIN, SATISH RAO, AND WARREN D. SMITH. Shallow excluded minors and improved graph decompositions. In DANIEL DOMINIC SLEATOR, ed., *Proc. 5th Annual ACM-SIAM Symp. on Discrete Algorithms (SODA '94)*, pp. 462–470. ACM, 1994. <http://dl.acm.org/citation.cfm?id=314464.314625>.
- [45] BRUCE A. REED. Tree width and tangles: a new connectivity measure and some applications. In *Surveys in combinatorics*, vol. 241 of *London Math. Soc. Lecture Note Ser.*, pp. 87–162. Cambridge Univ. Press, 1997. doi: 10.1017/CBO9780511662119.006. MR: 1477746.
- [46] BRUCE A. REED. Algorithmic aspects of tree width. In *Recent advances in algorithms and combinatorics*, vol. 11, pp. 85–107. Springer, 2003. doi: 10.1007/0-387-22444-0_4.
- [47] NEIL ROBERTSON AND PAUL SEYMOUR. Graph minors I–XXIII. *J. Combin. Theory Ser. B*, 1983–2010.
- [48] NEIL ROBERTSON AND PAUL SEYMOUR. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986. doi: 10.1016/0196-6774(86)90023-4. MR: 0855559.

- [49] NEIL ROBERTSON AND PAUL SEYMOUR. Graph minors. XVI. Excluding a non-planar graph. *J. Combin. Theory Ser. B*, 89(1):43–76, 2003. doi: 10.1016/S0095-8956(03)00042-X. MR: 1999736.
- [50] JAN VAN DEN HEUVEL, PATRICE OSSONA DE MENDEZ, DANIEL QUIROZ, ROMAN RABINOVICH, AND SEBASTIAN SIEBERTZ. On the generalised colouring numbers of graphs that exclude a fixed minor. *European J. Combin.*, 66:129–144, 2017. doi: 10.1016/j.ejc.2017.06.019.
- [51] DAVID R. WOOD. Bounded-degree graphs have arbitrarily large queue-number. *Discrete Math. Theor. Comput. Sci.*, 10(1):27–34, 2008. <http://dmtcs.episciences.org/434>. MR: 2369152.
- [52] DAVID R. WOOD. On tree-partition-width. *European J. Combin.*, 30(5):1245–1253, 2009. doi: 10.1016/j.ejc.2008.11.010. MR: 2514645.
- [53] DAVID R. WOOD. Clique minors in cartesian products of graphs. *New York J. Math.*, 17:627–682, 2011. <http://nyjm.albany.edu/j/2011/17-28.html>.
- [54] XUDING ZHU. Colouring graphs with bounded generalized colouring number. *Discrete Math.*, 309(18):5562–5568, 2009. doi: 10.1016/j.disc.2008.03.024.